



FIJIANATIONAL UNIVERSITY

**COLLEGE OF AGRICULTURE, FISHERIES & FORESTRY
SCHOOL OF AGRICULTURAL SCIENCE
DEPARTMENT OF AGRICULTURAL ECONOMICS AND
EXTENSION EDUCATION**

Bachelor of Science in Fisheries – Year II

**AGS 701: AGRICULTURAL STATISTICS, RESEARCH
METHODS AND EXPERIMENTAL DESIGNS**

FINAL EXAMINATION – TRIMESTER 3, 2016

*Time Allowed: 3 hours plus 10 minutes reading
Total marks: 100*

INSTRUCTIONS

1. This paper consists of **three** sections and **12** pages.
2. Answer all questions in the answer booklet provided.
3. Make sure to indicate your **identification number** in all pages you use.
4. You can use permitted calculators.
5. Statistical Tables are attached with list of formulae.
6. This exam is worth 50% of your overall mark.

SECTION A	20 MULTIPLE CHOICE QUESTIONS	20 MARKS
SECTION B	PART I: 10 TRUE/FALSE QUESTIONS PART II: 10 FILL IN THE BLANKS	10 MARKS 10 MARKS
SECTION C	5 LONG ANSWER QUESTIONS	60 MARKS

SECTION A:**MULTIPLE CHOICE****(20 MARKS)**

This section consists of 20 multiple choice questions worth 1 mark each. Write the letter corresponding to the best answer in the Answer Booklet provided.

1. The 'number of fish in Naduruloulou' the freshwater Fish Farm, is an example of which type of variable?

(A) a discrete variable	(B) a continuous variable
(C) a qualitative variable	(D) none of the above

2. In which of the following statement, an inferential statistics have been used?

(A) In the year 2010, 250 students were enrolled in CAFF Koronivia.
(B) 8 out 10 lecturers at FNU are men.
(C) Expenditures for FNU were \$10.1 million in 1999.
(D) Drinking decaffeinated coffee can raise cholesterol levels by 7%.

3. A researcher divided subjects into two groups according to breed of fish and then selected fish from each group for her sample. What sampling method was the researcher using?

(A) Cluster sampling	(B) Stratified sampling
(C) Simple Random sampling	(D) Systematic sampling

4. Which of the following is an appropriate measure of central tendency for nominal data?

(A) Mean.	(B) Median.	(C) Mode.	(D) Midrange.
-----------	-------------	-----------	---------------

5. The advantage of stem and leaf plots over grouped frequency distribution is that it

(A) is more reliable.
(B) can be used when there are lots of data values.
(C) is a more systematic way to organize data.
(D) can retain the actual data while showing them in graphical form.

6. What is the value of the mode when all values in the data set are different?


(A) 0
(B) 1
(C) There is no mode.
(D) It cannot be determined unless the data values are given.

7. Let X be the number of days per week that 30 AGS701 students do a 30 minute work on the Koronivia Farm.

X	Number of Students
0	3
1	2
2	3
3	8
4	1
5	9
6	4

The mean is:

- (A) 5 (B) 8 (C) 3.5 (D) 4
8. If the mode is to the left of the median and the mean is to the right of the median, then the distribution is:
- (A) Right skewed
(B) Left skewed
(C) Symmetrical
(D) Uniformed
9. When the value of α is increased, the probability of committing a type I error is
- (A) Decreased (B) Increased (C) The same (D) None of the above
10. Which type of alternative hypothesis is used in the figure below?



- (A) $H_1: \mu = k$ (B) $H_1: \mu \neq k$ (C) $H_1: \mu > k$ (D) $H_1: \mu < k$

11. If you wish to test the claim that the mean of the population is 100, the appropriate null hypothesis is:

- (A) $\bar{x} = 100$ (B) $\mu \leq 100$ (C) $\mu \geq 100$ (D) $\mu \neq 100$

12. For the t-test, one uses _____ instead of σ .

- (A) n (B) s (C) μ (D) \bar{x}


13. When the population standard deviation is unknown and the sample size is less than 30, what table value should be used in computing the test value for the hypothesis testing

- (A) z (B) t (C) N^2 (D) F

14. In regression, the variable being predicted is usually referred to as the

- (A) dependent variable (B) independent variable
(C) coefficient of correlation (D) coefficient of determination

15. The following scatter plot indicates



- (A) strong positive correlation (B) no correlation
(C) positive correlation (D) negative correlation

16. The formula for coefficient of variation of a sample is :

- (A) $CV = \frac{\mu}{\sigma}$ (B) $CV = \frac{\sigma}{\mu}$
(C) $CV = \frac{\bar{X}}{s}$ (D) $CV = \frac{s}{\bar{X}}$

17. In ANOVA, the null hypothesis should be rejected only

- (A) when there is no difference among all pairs of means.
(B) when there is no difference among all pairs of variances.
(C) when there is a difference among all pairs of means.
(D) none of the above.

18. Determining the table value for the F-distribution is different from finding values in the t-distribution tables because the F-table requires _____ value/s for degrees of freedom.
- (A) one (B) two (C) three (D) more than three
19. In experimental designs, a collection of plots is termed as
- (A) experimental unit (B) Plot
(C) block (D) treatment
20. Assigning of treatments or factors to be tested to the experimental units according to definite laws or probability is technically known as
- (A) Randomization (B) Replication
(C) Local Control (D) Experimentation

SECTION B:

(20 MARKS)

Part I:

True/False Questions

(10 marks)

In the Answer Booklet provided write true or false for the following questions.

1. The variable temperature is an example of a quantitative variable.
2. It is not important to keep the width of each class the same in a frequency distribution.
3. In construction of a frequency polygon, the class limits are used for the x-axis.
4. When the mean is computed for individual data, all values in the data set are used.
5. An outlier affects the median more than the mean.
6. The positive square root of the variance is called standard deviation.
7. No error is committed when the null hypothesis is rejected when it is false.
8. The test value separates the rejection region from the acceptance region.
9. A correlation coefficient of +1 implies a strong positive linear relationship between the variables.
10. A negative relationship between two variables means that for the most part, as the x variable increases, the y variable increases.

Part II:**Fill in the Blanks****(10 marks)**

Fill in the blanks with (word or phrase or symbol or letter) the appropriate answer in the Answer Booklet.

1. A group of plants selected from the group of all plants under study is called a _____.
2. The three types of frequency distributions are _____, Ungrouped and Grouped.
3. Picking every 10th fish from a large pool for study would be an example of _____ sampling.
4. Two major branches of statistics are Descriptive and _____.
5. A measure obtained from sample data is called a sample _____.
6. The symbol for population standard deviation is _____.
7. An extremely high or extremely low data value is called an _____.
8. To test the claim that the mean is greater than 87, you would use a _____-tailed test.
9. The range of 'r' is from _____ to _____.
10. The regression line is also called the _____.

SECTION C:**LONG ANSWER QUESTIONS****(60 MARKS)**

This section consists of 5 long answer questions worth 12 marks each. Write your answers in the Answer Booklet provided. Show all necessary working as partial marks will be awarded to partially correct answers.

QUESTION 1*Start on a new page***[4+2+4+2=12 marks]**

The following data give the marks of 21 AGS 701 students at FNU CAFF.

41	54	22	28	31	39	58
63	48	67	47	58	37	26
55	61	47	59	36	48	54

- (i) Construct a frequency distribution table. Take 5 classes.
- (ii) Calculate the relative frequencies and percentages for all classes.
- (iii) Construct a histogram and frequency polygon for the relative frequency distribution.
- (iv) Construct an Ogive.

QUESTION 2*Start on a new page***[2+2+1+2+1+1+2+1= 12 marks]**

A. A survey of 30 supermarkets reported these numbers of fish sales during a randomly selected year.

Number of Sales	8-12	13-17	18-22	23-27	28-32	33-37
Frequency	1	4	10	5	4	6

Calculate the following:

- (i) Mean
- (ii) Median
- (iii) Mode
- (iv) Variance
- (v) Standard Deviation
- (vi) Coefficient of Variation
- (vii) Mean Deviation
- (viii) Skewness

QUESTION 3*Start on a new page***[6+6 =12 marks]**

A. A study claims that all adults spend an average of 14 hours on chores during a weekend. A researcher wanted to check if this claim is true. A random sample of 200 adults taken by this researcher showed that these adults spend an average of 13.75 hours on chores during a weekend with a standard deviation of 3 hours.

Will you reject the null hypothesis at $\alpha = 0.05$?

B. A fish researcher claims that the average number of infections per week in a pond is 16.3. A random sample of 10 fish had a mean number of 17.7 infections. The sample standard deviation is 1.8. Is there enough evidence to reject the investigator's claim at $\alpha = 0.05$?

QUESTION 5*Start on a new page***[4+3+3+2 = 12 marks]**

An Agricultural Statistics student for a survey recorded the seed yield per plant (in grams) (y) and plant height (in centimeters) (x) of bean as shown in the following table:

Seed yield per plant (in g), y	5.22	8.13	6.52	4.16	8.98	3.05	3.49	5.40	2.39	2.71	3.97	7.56
Plant height (in cm), x	94.2	69.3	115.3	83.3	85.4	68.1	50.7	96.2	76.1	52.0	82.1	81.3

- (i) Compute the value of the correlation coefficient and interpret.
- (ii) Find the coefficient of determination and interpret what it means.
- (iii) Determine the regression line equation
- (iv) Predict seed yield per plant, y , when the plant height, x , is 100cm.

QUESTION 6*Start on a new page***[12 marks]**

A research group desired to compare the meat production of different species of fish from Naduruloulou Farm. Three different fish species (namely Kanace, Malea, Tilapia) meat production (in kg) was recorded as shown below. At 5% level of significance is there a sufficient evidence to conclude that a difference in meat production exists? Use One-way ANOVA.

Kanace	Malea	Tilapia
5.99	8.99	4.99
6.99	7.99	3.99
8.59	6.29	5.29
6.49	7.29	4.49

THE END

LIST OF FORMULAE:

$$1. \quad \bar{X} = \frac{\sum f \cdot X_m}{n} \quad \text{and} \quad S^2 = \frac{\sum f(X_m - \bar{X})^2}{n-1}$$

$$2. \quad MD = l + \frac{N/2 - m}{f} \times c$$

$$3. \quad r = \frac{n(\sum xy) - (\sum x)(\sum y)}{\sqrt{n(\sum x^2) - (\sum x)^2} \cdot \sqrt{n(\sum y^2) - (\sum y)^2}}$$

4. The regression line $y' = a + bx$,
where

$$a = \frac{(\sum y)(\sum x^2) - (\sum x)(\sum xy)}{n(\sum x^2) - (\sum x)^2}$$

$$b = \frac{n(\sum xy) - (\sum x)(\sum y)}{n(\sum x^2) - (\sum x)^2}$$

$$5. \quad z = \frac{\bar{X} - \mu}{\sigma / \sqrt{n}}$$

$$6. \quad t = \frac{\bar{X} - \mu}{s / \sqrt{n}}$$

$$7. \quad S_B^2 = \frac{\sum n_i (\bar{X}_i - \bar{X}_{GM})^2}{k-1},$$

$$S_W^2 = \frac{\sum (n_i - 1) S_i^2}{\sum (n_i - 1)},$$

$$F = \frac{S_B^2}{S_W^2}$$

TABLE 1: The t-Distribution Table

The entries in this table give the critical values of t for the specified number of degrees of freedom and areas in the right tail.




df	Area in the Right Tail under the t Distribution Curve					
	.10	.05	.025	.01	.005	.001
1	3.078	6.314	12.706	31.821	63.657	318.309
2	1.886	2.920	4.303	6.965	9.925	22.327
3	1.638	2.353	3.182	4.541	5.841	10.215
4	1.533	2.132	2.776	3.747	4.604	7.173
5	1.476	2.015	2.571	3.365	4.032	5.893
6	1.440	1.943	2.447	3.143	3.707	5.208
7	1.415	1.895	2.365	2.998	3.499	4.785
8	1.397	1.860	2.306	2.896	3.355	4.501
9	1.383	1.833	2.262	2.821	3.250	4.297
10	1.372	1.812	2.228	2.764	3.169	4.144
11	1.363	1.796	2.201	2.718	3.106	4.025
12	1.356	1.782	2.179	2.681	3.055	3.930
13	1.350	1.771	2.160	2.650	3.012	3.852
14	1.345	1.761	2.145	2.624	2.977	3.787
15	1.341	1.753	2.131	2.602	2.947	3.733
16	1.337	1.746	2.120	2.583	2.921	3.686
17	1.333	1.740	2.110	2.567	2.898	3.646
18	1.330	1.734	2.101	2.552	2.878	3.610
19	1.328	1.729	2.093	2.539	2.861	3.579
20	1.325	1.725	2.086	2.528	2.845	3.552
21	1.323	1.721	2.080	2.518	2.831	3.527
22	1.321	1.717	2.074	2.508	2.819	3.505
23	1.319	1.714	2.069	2.500	2.807	3.485
24	1.318	1.711	2.064	2.492	2.797	3.467
25	1.316	1.708	2.060	2.485	2.787	3.450
26	1.315	1.706	2.056	2.479	2.779	3.435
27	1.314	1.705	2.052	2.473	2.771	3.421
28	1.313	1.701	2.048	2.467	2.763	3.408
29	1.311	1.699	2.045	2.462	2.756	3.396
30	1.310	1.697	2.042	2.457	2.750	3.385
31	1.309	1.696	2.040	2.453	2.744	3.375
32	1.309	1.694	2.037	2.449	2.738	3.366
33	1.308	1.692	2.035	2.445	2.733	3.356
34	1.307	1.691	2.032	2.441	2.728	3.346
35	1.306	1.690	2.030	2.438	2.724	3.340

TABLE 2: Areas under Standard Normal Probability Curve (Source: Eton Table)

z											1			4			7		
	0	1	2	3	4	5	6	7	8	9	2	3	4	5	6	7	8	9	
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359	4	8	12	16	20	24	28	32	36
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0754	4	8	12	16	20	24	28	32	36
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141	4	8	12	15	19	22	27	31	35
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517	4	6	11	15	19	22	26	30	34
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879	4	7	11	14	18	22	25	29	32
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224	3	7	10	14	17	21	24	27	31
0.6	.2258	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2518	.2549	3	6	10	13	16	19	23	26	29
0.7	.2580	.2612	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852	3	6	9	12	15	18	21	24	27
0.8	.2881	.2910	.2939	.2967	.2996	.3023	.3051	.3078	.3106	.3133	3	6	8	11	14	17	19	22	25
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389	3	5	8	10	13	15	18	20	23
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621	2	5	7	9	12	14	16	18	21
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830	2	4	6	8	10	12	14	16	19
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015	2	4	5	7	9	11	13	15	16
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177	2	3	5	6	8	10	11	13	14
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319	1	3	4	6	7	8	10	11	13
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441	1	2	4	5	6	7	8	10	11
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545	1	2	3	4	5	6	7	8	9
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633	1	2	3	3	4	5	6	7	8
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706	1	1	2	3	4	4	5	6	6
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767	1	1	2	2	3	4	4	5	5
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817	0	1	1	2	2	3	3	4	4
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857	0	1	1	2	2	2	3	3	4
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890	0	1	1	1	2	2	2	3	3
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916	0	0	1	1	1	2	2	2	2
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936	0	0	1	1	1	1	1	2	2
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952	0	0	0	1	1	1	1	1	1
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964	0	0	0	0	1	1	1	1	1
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974	0	0	0	0	0	1	1	1	1
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981	0	0	0	0	0	0	0	0	1
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986	0	0	0	0	0	0	0	0	1
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990	0	0	0	0	0	0	0	0	0
3.1	.4990	.4991	.4991	.4991	.4992	.4992	.4992	.4992	.4993	.4993	0	0	0	0	0	0	0	0	0
3.2	.4993	.4993	.4994	.4994	.4994	.4994	.4994	.4995	.4995	.4995	0	0	0	0	0	0	0	0	0
3.3	.4995	.4995	.4995	.4996	.4996	.4996	.4996	.4996	.4996	.4997	0	0	0	0	0	0	0	0	0
3.4	.4997	.4997	.4997	.4997	.4997	.4997	.4997	.4997	.4997	.4998	0	0	0	0	0	0	0	0	0
3.5	.4998	.4998	.4998	.4998	.4998	.4998	.4998	.4998	.4998	.4998	0	0	0	0	0	0	0	0	0
3.6	.4998	.4998	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	0	0	0	0	0	0	0	0	0
3.7	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	0	0	0	0	0	0	0	0	0
3.8	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	.4999	0	0	0	0	0	0	0	0	0
3.9	.5000	.5000	.5000	.5000	.5000	.5000	.5000	.5000	.5000	.5000	0	0	0	0	0	0	0	0	0

TABLE 3: F-DISTRIBUTION



The tabulated value is the value of F with v_1 , v_2 degrees of freedom which is exceeded with a probability of 5%. For other notes see i& table
 e.g. $P(F_{15,7} > 3.5168) = 5\%$.

$v_2 \backslash v_1$	1	2	3	4	5	6	7	8	9	10	12	15	20	30	60	∞
1	161.45	199.50	215.71	224.58	230.16	233.99	236.77	238.88	240.54	241.88	243.91	245.95	248.01	250.09	252.20	254.32
2	18.513	19.000	19.164	19.247	19.296	19.330	19.353	19.371	19.385	19.396	19.413	19.429	19.446	19.462	19.479	19.496
3	10.128	9.5521	9.2766	9.1172	9.0135	8.9406	8.8868	8.8452	8.8123	8.7855	8.7446	8.7029	8.6602	8.6166	8.5720	8.5265
4	7.7086	6.9443	6.5914	6.3883	6.2560	6.1631	6.0942	6.0410	5.9988	5.9644	5.9117	5.8578	5.8025	5.7459	5.6878	5.6281
5	6.6079	5.7861	5.4095	5.1922	5.0503	4.9503	4.8759	4.8183	4.7725	4.7351	4.6777	4.6188	4.5581	4.4957	4.4314	4.3650
6	5.9874	5.1433	4.7571	4.5337	4.3874	4.2839	4.2066	4.1468	4.0990	4.0600	3.9999	3.9381	3.8742	3.8082	3.7398	3.6688
7	5.5914	4.7374	4.3468	4.1203	3.9715	3.8660	3.7870	3.7257	3.6767	3.6365	3.5747	3.5108	3.4445	3.3758	3.3043	3.2298
8	5.3177	4.4590	4.0662	3.8378	3.6875	3.5806	3.5005	3.4381	3.3881	3.3472	3.2840	3.2184	3.1503	3.0794	3.0053	2.9276
9	5.1174	4.2565	3.8626	3.6331	3.4817	3.3738	3.2927	3.2296	3.1789	3.1373	3.0729	3.0061	2.9365	2.8637	2.7872	2.7067
10	4.9646	4.1028	3.7083	3.4780	3.3258	3.2172	3.1355	3.0717	3.0204	2.9782	2.9130	2.8450	2.7740	2.6996	2.6211	2.5379
11	4.8443	3.9823	3.5874	3.3567	3.2039	3.0946	3.0123	2.9480	2.8962	2.8536	2.7876	2.7186	2.6464	2.5705	2.4901	2.4045
12	4.7472	3.8853	3.4903	3.2592	3.1059	2.9961	2.9134	2.8486	2.7964	2.7534	2.6866	2.6169	2.5436	2.4663	2.3842	2.2962
13	4.6672	3.8056	3.4105	3.1791	3.0254	2.9153	2.8321	2.7669	2.7144	2.6710	2.6037	2.5331	2.4589	2.3803	2.2966	2.2064
14	4.6001	3.7389	3.3439	3.1122	2.9582	2.8477	2.7642	2.6987	2.6458	2.6021	2.5342	2.4630	2.3879	2.3082	2.2230	2.1307
15	4.5431	3.6823	3.2874	3.0556	2.9013	2.7905	2.7066	2.6408	2.5876	2.5437	2.4753	2.4035	2.3275	2.2468	2.1601	2.0658
16	4.4940	3.6337	3.2389	3.0069	2.8524	2.7413	2.6572	2.5911	2.5377	2.4935	2.4247	2.3522	2.2756	2.1938	2.1058	2.0096
17	4.4513	3.5915	3.1968	2.9647	2.8100	2.6987	2.6143	2.5480	2.4943	2.4499	2.3807	2.3077	2.2304	2.1477	2.0584	1.9604
18	4.4139	3.5546	3.1599	2.9277	2.7729	2.6613	2.5767	2.5102	2.4563	2.4117	2.3421	2.2686	2.1906	2.1071	2.0166	1.9168
19	4.3808	3.5219	3.1274	2.8951	2.7401	2.6283	2.5435	2.4768	2.4227	2.3779	2.3080	2.2341	2.1555	2.0712	1.9796	1.8780
20	4.3513	3.4928	3.0984	2.8661	2.7109	2.5990	2.5140	2.4471	2.3928	2.3479	2.2776	2.2033	2.1242	2.0391	1.9464	1.8432
21	4.3248	3.4668	3.0725	2.8401	2.6848	2.5727	2.4876	2.4205	2.3661	2.3210	2.2504	2.1757	2.0960	2.0102	1.9165	1.8117
22	4.3009	3.4434	3.0491	2.8167	2.6613	2.5491	2.4638	2.3965	2.3419	2.2967	2.2258	2.1508	2.0707	1.9842	1.8895	1.7831
23	4.2793	3.4221	3.0280	2.7955	2.6400	2.5277	2.4422	2.3748	2.3201	2.2747	2.2036	2.1282	2.0476	1.9605	1.8649	1.7570
24	4.2597	3.4028	3.0088	2.7763	2.6207	2.5082	2.4226	2.3551	2.3002	2.2547	2.1834	2.1077	2.0267	1.9390	1.8424	1.7331
25	4.2417	3.3852	2.9912	2.7587	2.6030	2.4904	2.4047	2.3371	2.2821	2.2365	2.1649	2.0889	2.0075	1.9192	1.8217	1.7110
26	4.2252	3.3690	2.9751	2.7426	2.5868	2.4741	2.3883	2.3205	2.2655	2.2197	2.1479	2.0716	1.9898	1.9010	1.8027	1.6906
27	4.2100	3.3541	2.9604	2.7278	2.5719	2.4591	2.3732	2.3053	2.2501	2.2043	2.1323	2.0558	1.9736	1.8842	1.7851	1.6717
28	4.1960	3.3404	2.9467	2.7141	2.5581	2.4453	2.3593	2.2913	2.2360	2.1900	2.1179	2.0411	1.9586	1.8687	1.7689	1.6541
29	4.1830	3.3277	2.9340	2.7014	2.5454	2.4324	2.3463	2.2782	2.2229	2.1768	2.1045	2.0275	1.9446	1.8543	1.7537	1.6377
30	4.1709	3.3158	2.9223	2.6896	2.5336	2.4205	2.3343	2.2662	2.2107	2.1646	2.0921	2.0148	1.9317	1.8409	1.7396	1.6223
40	4.0848	3.2317	2.8387	2.6060	2.4495	2.3359	2.2490	2.1802	2.1240	2.0772	2.0035	1.9245	1.8389	1.7444	1.6373	1.5089
60	4.0012	3.1504	2.7581	2.5252	2.3683	2.2540	2.1665	2.0970	2.0401	1.9926	1.9174	1.8364	1.7480	1.6491	1.5343	1.3893
120	3.9201	3.0718	2.6802	2.4472	2.2900	2.1750	2.0867	2.0164	1.9588	1.9105	1.8337	1.7505	1.6587	1.5543	1.4290	1.2539
∞	3.8415	2.9957	2.6049	2.3719	2.2141	2.0986	2.0096	1.9391	1.8800	1.8317	1.7545	1.6695	1.5727	1.4643	1.3343	1.1439

F-DISTRIBUTION